

## Giant Condyloma Acuminatum of Buschke and Lowenstein

Lalya I<sup>1</sup>, Maghous A<sup>2</sup>,  
Marnouche E<sup>2</sup> and Mansouri H<sup>1</sup>

A 55-year-old man without HIV infection history, presented with neglected giant peri-anal condyloma acuminatum evolving for more than 10 years. Histologic findings revealed a condyloma acuminatum of Buschke and Lowenstein tumor without verrucous carcinoma. A patient was admitted for radiation therapy in our department. Clinical examination revealed a huge infected mass, cauliflower-like, infiltrating the left buttock and perineum (**Figure 1**). 50 Gy in 25 fractions were prescribed using parallel opposed fields (AP-Posterior Anterior [PA]) and a 1 cm bolus. The second image shows the immediate response to irradiation therapy (**Figure 2**). Unfortunately, the patient was lost to follow-up. The purpose of this image is to show the condyloma acuminatum radiosensitivity

- 1 Department of Radiotherapy, Mohamed V Military Teaching Hospital, Rabat, Morocco
- 2 Resident of Radiation Oncology, National Institute of Oncology, Rabat

**Corresponding author:** Abdelhak Maghous

✉ [issamlalya@yahoo.fr](mailto:issamlalya@yahoo.fr)

Resident of Radiation Oncology, National Institute of Oncology, Rabat

Tel: 00212661572770



Figure 1



Figure 2